

Front

FOSOMIN[®] (Fosfomycin)

Capsule & Suspension

فاسومین

COMPOSITION:

Fosomin Capsule 500 mg:

Each capsule contains:
Fosfomycin Calcium EP equivalent to
Fosfomycin 500 mg.

Product Specs.: CCL Pharmaceuticals

Fosomin Suspension 250 mg/5 ml:

Each 5 ml contains:
Fosfomycin Calcium EP equivalent to
Fosfomycin 250 mg.

Product Specs.: CCL Pharmaceuticals

MICROBIOLOGY:

Fosfomycin is bactericidal antibiotic. Following active uptake into the cell it is reported to interfere with first step synthesis of bacterial cell walls. It is active invitro and in vivo against a wide range of Gram-positive and Gram-negative bacteria including staphylococcus aureus, Streptococci, E. Coli, Proteus, Pseudomonas, Serratia, Salmonella, Shigella & Haemophilus. To a lesser degree it is active against Enterococci, Proteus rettgeri and morgani, Klebsiella, and Enterobacter.

INDICATIONS:

Capsule or suspension are indicated in the treatment of Acute, Subacute and Chronic, Non complicated urinary infections, infections of the digestive tract, preferably of enteric localization, and Dermatological infections.

DOSAGE AND ADMINISTRATION:

Oral Preparation:

Adult:

1 capsule (500 mg) every 6 hours or 2 capsules:
(1g) every 8 hours.

Children:

1-2 teaspoonful (250-500 mg) of Fosomin suspension every 8 hours.

Nursing Infant:

1 Teaspoonful every 6 - 8 hours. (upto one year old).

Premature & New Born Infants:

½ Teaspoon (125mg) every 6 hours.

RECONSTITUTION:

Preparation Of Suspension:

Hold the bottle upside down and tap the bottom until all the powder is loosened, add boiled and cooled water to make 60 ml oral suspension.

INCOMPATIBILITIES:

None have been reported.

SIDE EFFECTS:

Oral administration of FOSOMIN caused rarely, soft feces or

Back

diarrhea. Cutaneous rashes may appear which usually subsides with oral antihistamines without stopping the treatment.

TOXICITY:

None have been reported.

INSTRUCTIONS FOR CAPSULE:

- Store below 30°C.
- Protect from heat, sunlight & moisture.
- Keep out of the reach of children.
- To be sold on the prescription of a registered medical practitioner only.

INSTRUCTIONS FOR SUSPENSION:

- Store in a cool dry place.
- Protect from heat, sunlight & moisture.
- Shake well before use.
- Keep bottle tightly closed after use.
- Use within 14 days after reconstitution.
- Keep out of the reach of children.
- To be sold on the prescription of a registered medical practitioner only.

PRESENTATION:

FOSOMIN Capsule 500 mg : Pack of 1x10 capsules.
FOSOMIN Suspension 250 mg/5 ml : Pack of 60 ml.

ہدایات برائے کپسول :
30 درجہ سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔
گرمی، دھوپ اور نمی سے بچائیں۔
بچوں کی پہنچ سے دور رکھیں۔
صرف ڈاکٹر کے نسخہ پر فروخت کریں۔

ہدایات برائے سسپنشن :
شکاف اور بھٹی جگہ پر رکھیں۔
گرمی، دھوپ اور نمی سے بچائیں۔
استعمال سے پہلے بوتل کا اچھی طرح ہلائیں۔
استعمال کے بعد بوتل کا اچھی طرح بند کر کے رکھیں۔
سسپنشن بنانے کے بعد ان کے اندر استعمال کریں۔
بچوں کی پہنچ سے دور رکھیں۔
صرف ڈاکٹر کے نسخہ پر فروخت کریں۔

FOR FURTHER INFORMATION PLEASE CONTACT:

131-B
25034-0002-003-0000-0000



Manufactured by:
CCL Pharmaceuticals (Pvt.) Ltd.
62 Industrial Estate, Kot Lakhpat, Lahore, Pakistan.