

# pulmonol®

Cough  
Syrup

پلمونول®  
شیرپ

## COMPOSITION:

Each 5 ml contains:

Chlorphenamine maleate BP	5 mg.
Terpin hydrate USP	10 mg.
Potassium bicarbonate BP	0.1 mg.
Ammonium chloride BP	25 mg.
Menthol BP	1 mg.
Aminophylline Ph. Eur	32 mg.
Potassium guaiaicol sulfonate USP	5 mg.
Potassium citrate BP	0.1 mg.
Flavor & base	q.s.

**Product Specs.:** CCL Pharmaceuticals

## INDICATIONS:

For all types of coughs and congestion.

## CONTRAINDICATIONS

Hypersensitivity to any ingredients of this product.

## WARNING & PRECAUTIONS:

Do not exceed the recommended dosage because nervousness, dizziness and excitability may occur. Chlorpheniramine Maleate may cause drowsiness. Avoid taking with alcoholic drinks or other sedatives. Ask the doctor before taking this medication if you have glaucoma or problems urinating due to enlarged prostate. Caution is also advised in patients with liver or kidney disease.

May cause drowsiness hence caution is advised if driving or operating machinery.

## PREGNANCY & LACTATION:

This product should be used with caution in pregnancy, particularly during the third trimester. Chlorpheniramine Maleate may pass into breast milk. It should be avoided by breastfeeding mothers.

## DRUG INTERACTIONS:

The actions of Chlorpheniramine Maleate may be prolonged and intensified by the concomitant use of monoamine oxidase inhibitors (MAOI's). In case of Ammonium chloride, the urine acidification may increase the renal excretion of quinidine. The aminoglycosides and erythromycin are more effective in an alkaline medium; urine acidification may diminish these drugs effectiveness in treating bacterial urinary tract infections.

## ADVERSE EFFECTS:

Drowsiness, dry mouth, nervousness, dizziness, headache, alterations in blood pressure and heart rate and gastrointestinal distress.

## DOSAGE:

**ADULTS:** 1 to 2 teaspoonful.

**CHILDREN:** ½ to 1 teaspoonful.

The dose to be mixed with equal amount of warm water and repeat 3 to 4 times a day or as directed by the physician.

## CAUTION:

Temperature variation may cause Turbidity / precipitation / thickening.

## INSTRUCTIONS:

- For oral use only.
- Close cap tightly after use.
- Keep out of the reach of children.
- To be sold on the prescription of a registered medical practitioner only.

## STORAGE:

- Store at cool and dry place (15-30°C).
- Do not freeze.
- Protect from heat, sunlight and moisture.

## SHAKE WELL BEFORE USE

**PRESENTATION** : Bottles of 120 ml and 450 ml.

طریقہ استعمال: بالغ: ایک سے دو چائے کے کچھ

بچے: نصف سے ایک چائے کچھ

خوراک: نیم گرم مسامی پانی ملا کر دن میں ۳ سے ہمار

یاد آگزی داریت کے مطابق استعمال کریں۔

استعمال سے قبل بوتل کو اچھی طرح ہلائیں۔

احتیاط: دوجہ حرارت میں تہہ ٹی سیرپ

کو ٹیفب یا ماسکتی ہے۔

داریات: صرف پیچہ ٹیلے استعمال کریں۔

استعمال کے بعد ڈسکن کا اچھی طرح بند کر دیں۔

بچوں کی کھینچ سے دور رکھیں۔

صرف مستند ڈاکٹر کے نسخہ پر فروخت کریں۔

استونج:

ٹھنڈی اور ٹنگل جلا (۱۵-۱۰ سینٹی گریڈ دوجہ حرارت) پر رکھیں۔

مجمد ہونے سے بچائیں۔ گرمی، دھوپ اور نمی سے بچائیں۔

Distributed by:

Neopharm Healthcare (Pvt.) Ltd.

65 Industrial Estate, Kot Lakhpat, Lahore, Pakistan.

FOR FURTHER INFORMATION PLEASE CONTACT:

 **CCL**  
Consumer Healthcare

Manufactured by:

CCL Pharmaceuticals (Pvt.) Ltd.

62 Industrial Estate, Kot Lakhpat, Lahore, Pakistan.

2845-A