

Herbigest

Syrup

Helps in

Heartburn, Indigestion
Abdominal Pain, Cramps,
Bloating Gas & Constipation

هربی گیسٹ
سیرپ

COMPOSITION:

Each 5 ml contains:

- *Iberis amara* (bitter candy tuft) whole plant ext. 15 mg (5:1) equiv. to raw herb (MS) 75 mg.
- *Angelica root extract* 3.33 mg (10:1) equiv. root dry (MS) 33.3 mg.
- *Matricaria chamomilla* (Chamomile) flower ext. 6.67 mg (10:1) equiv. flower dry (MS) 66.7 mg.
- *Carum carvi* (Caraway) herb ext. 8.325 mg (4:1) equiv. herb dry (MS) 33.3 mg.
- *Silybum marianum* (St. Mary's thistle) fruit ext. 3.33 mg (10:1) equiv. fruit dry (MS) 33.3 mg.
- *Melissa officinalis* (Lemon balm) leaf ext. 6.66 mg (5:1) equiv. leaf dry (MS) 33.3 mg.
- *Mentha x piperita* (Peppermint) leaf ext. 1.67 mg (10:1) equiv. leaf dry (MS) 16.7 mg.
- *Chelidonium majus* (Greater celandine) herb ext. 8.325 mg (5:1) equiv. herb dry (MS) 33.3 mg.
- *Glycyrrhiza glabra* (Liquorice) root ext. 3.33 mg (10:1) equiv. root dry (MS) 33.3 mg.

Product Specs.: In-house

How it Works?

The combination of 9 herbal extracts in **Herbigest** has been shown in controlled clinical trials to relieve gastric and abdominal discomfort associated with functional and motility-conditioned gastrointestinal disturbances such as functional dyspepsia and irritable bowel syndrome. This medicine is not intended to diagnose, treat, cure or prevent any disease.

INDICATIONS:

Herbigest contains medicinal plant extracts that work together to provide relief from multiple digestive symptoms listed below:

- Stomach pain
- Abdominal cramps
- Bloating
- Gas
- Heartburn
- Fullness
- Nausea
- Constipation
- Indigestion

DOSAGE AND ADMINISTRATION:

Adults and children over 12 years: Take 1 teaspoon (5.0 ml) 3 times a day.

Children 6 to 12 years: Give 3/4" of teaspoon (3.75 ml) 3 times a day.

Children 3 to 5 years: Give half teaspoon (2.5 ml) 3 times a day.

"SHAKE WELL BEFORE USE."

Herbigest should be taken before or with meals, three times a day. May be taken in some liquid (eg. warm water or any other choice of drink). The duration of use depends on functional digestive symptoms and how long they last for. Use for prolonged period should be done after consultation with a healthcare professional. If your symptoms persist then seek help from a healthcare professional.

CONTRAINDICATIONS:

Hypersensitivity to any of the ingredients of **Herbigest**.

WARNINGS AND SPECIAL PRECAUTIONS:

Patients with pre-existing liver disease should consult with their doctor before starting treatment on **Herbigest**.

Patients affected with serious chronic diseases and/or undergoing medical treatment should take **Herbigest** under medical guidance.

Herbigest may be taken by diabetics.

Do not give to children under 3 years.

INTERACTIONS:

There are no known interactions with other medicines.

PREGNANCY & LACTATION:

The use of **Herbigest** during pregnancy and lactation has not been fully studied and it should only be used in these circumstances under the guidance of a healthcare professional.

SIDE EFFECTS:

During clinical trials and routine use of no specific side effects have been reported.

OVERDOSAGE:

No reports of overdosage have been found in the literature. In cases of suspected overdosage treatment should be symptomatic and supportive.

INSTRUCTIONS:

- Store below 30°C.
- Protect from heat, sunlight & moisture.
- Keep out of the reach of children.
- Do not Refrigerate.

PRESENTATION:

Herbigest Syrup : Bottle of 60 ml syrup.

خوراک:

بالغ اور ۱۲ سال سے زائد عمر کے بچے: ایک چائے کاج (5.0 ملی لیٹر) دن میں ۳ بار لیں۔

۶ سے ۱۲ سال کی عمر کے بچے: تین چوتھائی چائے کاج (3.75 ملی لیٹر) دن میں ۳ بار لیں۔

۳ سے ۵ سال کی عمر کے بچے: آدھا چائے کاج (2.5 ملی لیٹر) دن میں ۳ بار لیں۔

یا ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

“استعمال سے پہلے اچھی طرح ہلائیں۔“

ہدایات:

• درجہ بندی گریڈ سے کم درجہ حرارت پر رکھیں۔

گرمی دھوپ اور نمی سے بچائیں۔

بچوں کی پہنچ سے دور رکھیں۔ فریج میں نہ رکھیں۔

Manufactured by:
Phytocon International (Pvt.) Ltd.
339-A, Sunder Industrial Estate, Raiwind Road,
Lahore, Pakistan.

Distributed by:
Nexpharm Healthcare (Pvt.) Ltd.

FOR FURTHER INFORMATION PLEASE CONTACT:



Manufactured for:
CCL Pharmaceuticals (Pvt.) Ltd.
62 Industrial Estate, Kot Lakhpat, Lahore, Pakistan.